

Heavy-flavour productions in the relativistic heavy ion collisions at the LHC

Shingo Sakai, Univ. of Tsukuba

Heavy Flavour (HF) in pp, p-Pb & Pb-Pb

- ▶ **Heavy-flavour (charm & beauty) production**
 - ▶ Initial hard scatterings ($M_{\text{HF}} \gg \Lambda_{\text{QCD}}$)
- ▶ **pp collisions**
 - ▶ Test for perturbative QCD (pQCD)
 - ▶ Reference for heavy ion collisions (both experiment & theory)
- ▶ **Heavy ion collisions**
 - ▶ Created in initial parton-parton scatterings
 - ▶ Traverse and interact with the hot & dense QCD matter
 - ▶ A good probe to study properties of the QCD matter
 - ▶ Energy loss (R_{AA}), collectivity (v_2), hadronization
- ▶ **pA collisions**
 - ▶ Control measurement for heavy ion collisions to disentangle initial from final state effects
 - ▶ Cold nuclear matter effect on heavy-flavour production

Energy Loss of heavy flavours

- In-medium parton energy loss
 - Radiative energy loss (PLB 632, 81)
 - gluon bremsstrahlung
 - smaller energy loss for heavy than for light quarks due to “dead cone” effect (PLB 519 (2001) 199.)
 - energy loss depends on the colour charge and is larger for gluons than for quarks
 - Collisional energy loss (PLB 649, 139)
 - energy loss via elastic scattering
- Theoretical predictions:
 - mass & colour charge dependence of energy loss
 - $E_{\text{loss}}(g) > E_{\text{loss}}(u,d,s) > E_{\text{loss}}(c) > E_{\text{loss}}(b)$

Nuclear modification factor

$$R_{AA}^{\pi} < R_{AA}^D < R_{AA}^B \quad ?$$

$$R_{AA}(p_T) = \frac{d N_{AA}/dp_T}{\langle T_{AA} \rangle \times d\sigma_{pp}/dp_T}$$

Azimuthal anisotropy of Heavy flavours

■ Elliptic flow

$$dN/d(\varphi-\psi_{RP}) = \dots + N_0(1+2v_2\cos(2(\varphi-\psi_{RP}))) + \dots$$

- Transfer initial spatial anisotropy to momentum anisotropy

- macroscopic: hydro model

- => pressure gradient

- microscopic

- => scattering in the medium

- Low p_T


- coupling of heavy quarks with the medium and their thermalization

- Intermediate p_T


- Hadronization mechanism (recombination)

- High p_T

- Path-length dependence of energy loss



Initial spatial anisotropy




Momentum space anisotropy of particle emission

▶ ***Heavy-flavour results in pp collisions***


Charm production in pp collisions

JHEP 1201 (2012) 128




ALI-PUB-111909

CMS-HIN-16-005





JHEP1603(2016)159





- D meson production mid- and forward-rapidity is in good agreement with pQCD calculations
 - upper side of the FONLL uncertainty band
 - various energies: 5.02, 7 and 13 TeV
 - from $p_T = 0$ to 100 GeV/c

Beauty production in pp collisions



- B meson production is in good agreement with pQCD calculations
 - FONLL is better agreement from low p_T to high p_T
 - PYTHIA overestimates at lower p_T
- b-jet production is also well represented by a pQCD (NLO)

HF production in pp collisions



- Productions of leptons (e, μ) from charm + beauty decays in different rapidity ranges are also well described by pQCD calculations

▶ ***Heavy-flavour results in p-Pb collisions***

p-A collisions

- ▶ Heavy-flavour in p-A collisions
 - ▶ control measurement for heavy-ion collisions to disentangle initial (cold nuclear matter effects) from final state effects

- ▶ Cold nuclear matter effects


- ▶ nuclear modification of Parton distribution Functions (PDF): shadowing or gluon saturation

K.J. Eskola et al., JHEP 0904(2009)65

H. Fuji & K. Watanabe, NPA 915 (2013) 1

- ▶ energy loss I. Vitev et al., PRC 75(2007) 064906
 - ▶ k_T broadening (Cronin enhancement)
 - ▶ multiple collisions


A.M. Glenn et al., PLB 644(2007)119



$$R_{pPb}(p_T) = \frac{d N_{pPb}/dp_T}{\langle T_{AA} \rangle \times d\sigma_{pp}/dp_T}$$

R_{pPb} of D, B and e^{HF} at mid-rapidity


PRL 113 (2014) 232301




ALI-PUB-79415

PRL 115 (2016) 032301

34.6 nb⁻¹ (pPb 5.02 TeV)



Phys. Lett. B 754 (2016) 81-93




ALI-PUB-100497


- R_{pPb} of D mesons, B mesons and e^{HF} is consistent with unity
 - No significant cold nuclear matter effects on heavy-flavour production
- Theoretical calculations with CNM effects are consistent with data
 - predict a small suppression at low p_T due to gluon saturation at low x
- Possible enhancement due to radial flow is predicted small based on Blast-wave model [PLB 731 (2014) 51]

R_{pPb} of c-jets and b-jets at mid-rapidity

CMS-HIN-15-012






PLB 754 (2016) 59



- Measured c-jet cross section in p-Pb is consistent with PYTHIA simulation
- R_{pPb} of b-jet with PYTHIA-based estimation is consistent with unity
 - considering the uncertainty on the PYTHIA reference

D production at forward-backward rapidity



LHCb-CO NF-2016-003



- D⁰ production at forward and backward rapidity
 - forward: p-going, $1.5 < y < -4$
 - backward: Pb-going, $-5 < y < -2.5$
- Significant D⁰ production asymmetry in forward – backward rapidity regions
- Measurements are consistent with a theoretical calculation
 - NLO with CTEQM and EPS09NLO

B- \rightarrow J/ Ψ production at forward-backward rapidity



JHEP 02 (2014) 072



- B- \rightarrow J/ Ψ production at $1.5 < \eta < 4.0$ (forward) and $-5 < \bar{\eta} < -2.5$ (backward)
- R_{FB} of B- \rightarrow J/ Ψ is asymmetry
 - backward yield is suppressed w.r.t. forward yield
- R_{FB} of B- \rightarrow J/ Ψ is larger than R_{FB} of prompt J/ Ψ
 - indicate cold nuclear matter effect is less pronounced for b hadrons


▶ ***Heavy-flavour results in Pb-Pb collisions***

D mesons in central Pb-Pb collisions




- Strong suppression of D mesons production
 - similar magnitude of suppression in 2.76 and 5.02 TeV
 - suppression observed up to 100 GeV/c at 5.02 TeV
 - D_s tends to larger : a hint of recombination process
- Suggest significant energy loss of charm in the medium

e^{HF} & μ^{HF} production in Pb-Pb collisions





ALI-PREL-101085



- Strong suppression of e^{HF} ($|y| < 0.6$) & μ^{HF} ($2.5 < y < 4$) in central collisions
 - similar suppression of e^{HF} & μ^{HF} in different rapidity regions
 - less suppression in mid-central collisions in both rapidity regions
 - high p_{T} : large contribution from beauty
- Suggest significant energy loss of charm and beauty in the medium

R_{AA} of B meson decays ($B \rightarrow e$ & $B \rightarrow J/\psi$)

JHEP 1205 (2012) 063




ALI-PREL-74678


- Suppression of $B \rightarrow e$ and $B \rightarrow J/\psi$ at high p_T
 - lower p_T : tends to follow binary scaling (consistent with unity)
 - high p_T (> 3 GeV/c): $R_{AA} \sim 0.4-0.5$
- Suggestions of beauty energy loss in the dense QCD matter

R_{AA} of charged particles, D and B \rightarrow J/ Ψ

CMS-PAS-HIN-16-001





JHEP 1511 (2015) 205, CMS-PAS-HN-12-014



- The magnitude of D meson suppression is similar to charged particles (π) within uncertainties
 - can't conclude on the expectation : $R_{AA} (D) > R_{AA} (\pi)$
- R_{AA} of D meson is smaller than R_{AA} of B \rightarrow J/ Ψ
 - indication of smaller energy loss of beauty than charm



Beauty jet

PRL 113 (2014) 132301




- Heavy-flavour jets: allow to address energy loss at parton level
- **Observed strong suppression of b-jets in most-central collisions**
 - similar magnitude of suppression to inclusive jet
 - **high p_T b-jets: largely comes from gluon splitting**

Imbalance of pairs of b jets




- Sub-leading recoil jets
 - larger path-length, primary b-jets from flavour creation
- Toward increasing imbalance with increasing centrality
 - similar imbalance as inclusive dijet



Azimuthal anisotropy of D mesons

PRC 90 (2014) 034904




ALI-PUB-70100

- Non zero D v_2 at low p_T
- Tends to get large from central (0-10%) to mid-central (30-50%)
 - Hydrodynamical behavior
- Consistent with charged particle v_2
- Charm quarks participate to the collective motion of the system

Azimuthal anisotropy of e^{HF} and μ^{HF}




e^{HF} : arXiv: 1606.00321, μ^{HF} : PLB 753 (2016) 41-56



- Non-zero v_2 of e^{HF} at $|y| < 0.7$ and μ^{HF} at $2.5 < y < 4$
 - the magnitude is compatible in mid- and forward-rapidities
- v_2 of e^{HF} measured from $p_T > 0.5$ GeV/c
 - similar p_T dependence to other light hadron v_2
- v_2 at high p_T e^{HF} and μ^{HF} reflects beauty
- Charm quarks participate to the collective motion of the system

Comparison with models (I)

JHEP09(2012)112





Theoretical calculations

- **initial:** with/without cold nuclear matter from PDF
- **medium modeling:** Hydro, Glauber, parton transportation
- **interaction:** radiative, collisional, resonant interaction
- **hadronization:** fragmentation, coalescence
- **Models represent R_{AA} of D mesons, e^{HF} and μ^{HF}**
 - mid- and forward-rapidity regions
 - high p_T leptons (e, μ) mainly from beauty decay

BAMPS: J. Phys. G 38 (2011) 124152,
POWLANG: Eur. Phys. J C
 71(2011)1666,
UrQMD: arXiv:1211.6912, J. Phys. Conf.
 Ser. 426,012032(2013),
TAMU: Phys. Rev. C 86 (2012) 014903,
WHDG: J. Phys. G38(2011)124114,
Aichelin: Phys. Rev. C79(2009)044906,
 J. Phys. G37(2010)094019
Cao, Qin, Bass: arXiv:1308.0617


Comparison with models (2)



Theoretical calculations

- **initial**: with/without cold nuclear matter from PDF
- **medium modeling**: Hydro, Glauber, parton transportation
- **interaction**: radiative, collisional, resonant interaction
- **hadronization**: fragmentation, coalescence
- Large suppression and non-zero v_2 (at low p_T) are represented by models, but simultaneous reproduction of the R_{AA} and v_2 is challenging

Comparison with models (3)



Experimental result

- $R_{AA}(D) < R_{AA}(B \rightarrow J/\Psi)$

Theoretical model

- radiative + collisional energy loss
- used two masses (charm and beauty) for calculating $B \rightarrow J/\Psi$ R_{AA} to study mass dependence
- result using beauty mass well represents centrality dependence of $R_{AA}(B \rightarrow J/\Psi)$
- the difference between D meson and $B \rightarrow J/\Psi$ is mainly from mass in this model

Summary

- ▶ **Heavy-flavour measurements at LHC**
 - ▶ D, B, leptons from heavy flavours, c-jet and b-jet
 - ▶ The productions are well described by pQCD calculations in pp collisions
- ▶ **Pb-Pb collisions**
 - ▶ **Strong suppression of heavy-flavour yield**
 - ▶ Clear indication for substantial energy loss of charm and beauty in the hot and dense matter
 - Not observed such suppression in pPb
 - ▶ Results indicate beauty lose smaller energy than charm
 - ▶ **Non-zero & centrality dependence of v_2**
 - ▶ Suggest strong re-interaction in the medium
- ▶ **Heavy flavours observed to be significantly affected by hot and dense QCD medium**